



RT-PCR and sequence analysis of *Macrobrachium rosenbergii* nodavirus: Indian isolate

M.S. Shekhar^{*}, I.S. Azad, K.P. Jithendran

Central Institute of Brackishwater Aquaculture, 75 Santhome High Road, R.A.Puram, Chennai-600 028, India

Received 20 September 2004; received in revised form 5 July 2005; accepted 25 July 2005

Abstract

White tail disease is a recently reported disease of giant freshwater prawn, *Macrobrachium rosenbergii* associated with *M. rosenbergii* nodavirus (*MrNV*). The present study is the first report on RT-PCR and sequence analysis of *MrNV* of an Indian isolate confirming the nodavirus infection in *M. rosenbergii* from India. Total RNA was extracted and the cDNA was synthesized from the infected post larvae collected from a hatchery located in Chennai, India. An 850-bp amplified product could be obtained by RT-PCR. The nucleotide sequence analysis of this 850-bp segment showed 98% nucleotide and 99% amino acid sequence identity with the reported sequence of an *MrNV* isolate from the West Indies.

© 2005 Elsevier B.V. All rights reserved.

Keywords: *Macrobrachium rosenbergii* nodavirus; RT-PCR; Sequence analysis

1. Introduction

Occurrence of white tail disease (WTD) in hatchery-reared *Macrobrachium rosenbergii* was first reported from the West Indies (Arcier et al., 1999). The proposed causative agent was identified as an icosahedral, non-enveloped virus of 26–27 nm called *M. rosenbergii* nodavirus (*MrNV*). The genome of this virus contains two pieces of ssRNA, of 3.0 kb (RNA-1) and 1.2 kb (RNA-2) (Qian et al., 2003). Along with this virus, extra small virus

particles (XSV) of 14–16 nm have been found in whitish muscle disease cases in prawns from China. XSV has been proposed to be a satellite virus dependent on the larger *MrNV* particles (Qian et al., 2003). Actual role of XSV and its interactions with *MrNV* in WTD is still unclear as some positively diagnosed *MrNV* samples were found negative for XSV (Widada et al., 2004). In addition to *MrNV*, two other viruses, parvo like virus (Anderson et al., 1990) and *Macrobrachium* muscle virus (Tung et al., 1999) have been reported to affect this species. The occurrence of WTD has also been reported from Taiwan (Tung et al., 1999) and China (Qian et al., 2003), but there are no published reports of it from any other part of the world.

^{*} Corresponding author.

E-mail address: msshkhar@hotmail.com (M.S. Shekhar).




Fig. 1. RT-PCR product of *MrNV* (Indian isolate). Lane 1: 100 bp marker. Lane 2: PCR product of 850 bp obtained by RT-PCR.

Mortalities due to WTD in *M. rosenbergii* have been recently noticed in India from hatcheries located at Nellore, Andrapradesh and Chennai, Tamil Nadu. The present work is the first report confirming the presence of *MrNV* in *M. rosenbergii* in India by reverse transcriptase-polymerase chain reaction (RT-PCR) and sequencing of the PCR-amplified product.

2. Materials and methods

2.1. Samples

Post larvae (PL) of *M. rosenbergii* were obtained from a hatchery in Chennai, India. Mortality was recorded in the 8- to 10-day-old PL showing empty guts and whitish opaque tails.

2.2. RT-PCR

Total RNA was extracted from the infected PL using a NucleoSpin RNA II kit (Macherey-Nagel). The cDNA was synthesized and RT-PCR was carried out using a

```

1 atggaagtccgocgattaattgaagcatttgtgaagaatgaaccg
  M E V R R L I E A F V K N E P      15
46 accaataaatctggctgcataataatcgtctttcgcgactcaaga
  T N K S G R I I S S F A D S R      30
91 ttcttgttgaagttttccacatatacgccttgcccttcgagatgaa
  F L L K F S T Y T L A F R D E      45
136 gtattacatgctgaacataatcgacattggttttgtcctggattg
  V L H A E H N R H W F C P G L      60
181 acacctaaatgagatagcagataaagtctgcgactatgtctcgtggt
  T P N E I A D K V C D Y V R G      75
226 gttgcgacacctgcagaaggtgatttttagcaactttgacggaagg
  V A T P A E G D F S N F D G R      90
271 gtatctgcttgggtgtcaagagaacgtgatgaatgcggtttaccac
  V S A W C Q E N V M N A V Y H      105
316 agatgggtttaaccgtaagttttcttaaggaattgcagaagtataca
  R W F N R K F S K E L Q K Y T      120
361 tcaatgttggtagttgcccagctcgagctaagcgttttggtttc
  S M L V S C P A R A K R F G F      135
406 cagtatgaaccgggagtggtgggttaagatggttagtccaaccacc
  Q Y E P G V G V K S G S P T T      150
451 tgtgaccttaattcagttctaaataacttactcaatacgcagca
  C D L N S V L N N F T Q Y A A      165
496 gttaggctgactaaaccagacctctcaccacaagaagcctttgaa
  V R L T K P D L S P Q E A F E      180
541 caaactggcttaagtttcggcgacgattcactatttgacaagcaa
  Q T G L S F G D D S L F D K Q      195
586 taccagctcagatggaattacgtcgtcgaacaacttggtatggaa
  Y Q L R W N Y V V E Q L G M E      210
631 ctcaaggttgaacccttcgacccaataacggtgtgacttttctt
  L K V E P F D P N N G V T F L      225
676 gctcgtgttttctctgatccttatagtacaaatactagttttcag
  A R V F P D P Y S T N T S F Q      240
721 gatccactaagaacgtgg
  D P L R T W

```

Fig. 2. Sequence of Indian *MrNV*. The numbering for the nucleotide sequence is given on left side and the numbering for the corresponding amino acid residues are given on the right side.

RobusT II RT-PCR kit (Finnzymes). RT was carried out at 42 °C for 60 min. The PCR cycle consisted of initial denaturation at 94 °C for 2 min followed by 30 cycles of denaturation at 92 °C for 1 min, annealing at 55 °C for 1 min and extension at 72 °C for 1 min with a final cycle of 10 min extension at 72 °C. The upstream primer 5'CCACGTTCTTAGTGGATCCT 3' and the downstream primer 5'CGTCCGCTGGTAGTTCC 3' (30 pmol each) specific to RNA-1 were used as reported by Widada et al. (2003). The PCR product was purified

by gel extraction using a NucleoSpin Extract kit (Macherey-Nagel).

2.3. Sequencing and sequence analysis

Sequencing of the gel-purified PCR product was carried out using both upstream and downstream primers by Bangalore Genei, India. The sequence was aligned and edited using BioEdit Sequence Alignment Editor. Sequence analysis was done using BLAST

<i>Epinephelus</i> NNV	DEFTMMVKAFQKKEAYNAPNYPRNISTVPHT-QNVKLSSTYAFKASVLQHVP---WYMP	56
Sevenband grouper NNV	DEFTMMVKAFQKKEAYNAPNYPRNISTVPHT-QNVKLSSTYAFKASVLQHVP---WYMP	56
Redspotted grouper NNV	DEFTMMVKAFQKKEAYNAPNYPRNISTVPHT-QNVKLSSTYAFKASVLQHVP---WYMP	56
Japanese flounder NNV	GEFTMIVKAFQKKEAYNAPNYPRNISTVPHT-QNVKLSSTYAFKASVLQHVP---WYMP	56
Atlantic halibut NNV	DEFTMIVKAFQKKEAYNAPNYPRNISTVPHT-QNVKLSSTYAFKASVLQHVP---WYMP	56
Striped jack NNV	DEFTMIVKAFQKKEAYNAPNYPRNISTVPHT-QNVKLSSTYAFKASVLQHVP---WYMP	56
<i>MrNV</i> (West Indies)	MEVRRLEIAFVKNEPTNKS--GRIISSFADSRFLKFKSTYTLAFREVLHAEHNRWFCP	58
<i>MrNV</i> (China)	MEVRRLEIAFVKNEPTNKS--GRIISSFADSRFLKFKSTYTLAFREVLHAEHNRWFCP	58
<i>MrNV</i> (India)	MEVRRLEIAFVKNEPTNKS--GRIISSFADSRFLKFKSTYTLAFREVLHAEHNRWFCP	58
	* . : : * * * * . * . * * * : : * : : * * * * : * : * : *	
<i>Epinephelus</i> NNV	THTPAEIAQAVNLAASSTELVETDYSKFDGTFLLRFMRCEVEFAIYKRWVHLDHLP	116
Sevenband grouper NNV	THTPAEIAQAVNLAASSTELVETDYSKFDGTFLLRFMRCEVEFAIYKRWVHLDHLP	116
Redspotted grouper NNV	THTPAEIAQAVNLAASSTELVETDYSKFDGTFLLRFMRCEVEFAIYKRWVHLDHLP	116
Japanese flounder NNV	THTPAEIAQAVNLAASSTELVETDYSKFDGTFLLRFMRCEVEFAIYKRWVHLDHLP	116
Atlantic halibut NNV	THTPAEIAQAVNLAASSTELVETDYSKFDGTFLLRFMRCEVEFAIYKRWVHLDHLP	116
Striped jack NNV	THTPAEIAQAVNLAASSTELVETDYSKFDGTFLLRFMRCEVEFAIYKRWVHLDHLP	116
<i>MrNV</i> (West Indies)	GLTPNEIADKVCQYVVRGATPAEGDFSNFDGRVSAWQENVMNAVYHRWFNRKFSKELQ	118
<i>MrNV</i> (China)	GLTPNEIADKVCQYVVRGATPAEGDFSNFDGRVSAWQENVMNAVYHRWFNRKFSKELQ	118
<i>MrNV</i> (India)	GLTPNEIADKVCQYVVRGATPAEGDFSNFDGRVSAWQENVMNAVYHRWFNRKFSKELQ	118
	** * * * : * : * * * * : * : : * * * * : * : * : *	
<i>Epinephelus</i> NNV	LLANEIQAPAVTR-LGIKYDPPDCSRLSGSALTTDGNISIANAFVSYLAGRMAG--MDDDEA	173
Sevenband grouper NNV	LLANEIQAPAVTR-LGIKYDPPDCSRLSGSALTTDGNISIANAFVSYLAGRMAG--MDDDEA	173
Redspotted grouper NNV	LLANEIQAPAVTR-LGIKYDPPDCSRLSGSALTTDGNISIANAFVSYLAGRMAG--MDDDEA	173
Japanese flounder NNV	LLANEIQAPAVTR-LGIKYDPPDCSRLSGSALTTDGNISIANAFVSYLAGRMAG--MDDDEA	173
Atlantic halibut NNV	LLANEIQAPAVTR-LGIKYDPPDCSRLSGSALTTDGNISIANAFVSYLAGRMAG--MDDDEA	173
Striped jack NNV	LLANEIQAPAVTR-LGIKYDPPDCSRLSGSALTTDGNISIANAFVSYLAGRMAG--MDDDEA	173
<i>MrNV</i> (West Indies)	YTSMLVSCPARAKRFGFQYEPGVGVKSGSPTTCDLNSVLNFTQYAAVRLTKPDLSPQEA	178
<i>MrNV</i> (China)	YTSMLVSCPARAKRFGFQYEPGVGVKSGSPTTCDLNSVLNFTQYAAVRLTKPDLSPQEA	178
<i>MrNV</i> (India)	YTSMLVSCPARAKRFGFQYEPGVGVKSGSPTTCDLNSVLNFTQYAAVRLTKPDLSPQEA	178
	: : : * : : * * * * : * * * * : * * * * : * * * * : * * * * : * * * * :	
<i>Epinephelus</i> NNV	WSWIGIVYGDGLRSGNVSNELLTNTASSLGFDLKIVNRAPRGSPTVFLSRVYLDPWSSP	233
Sevenband grouper NNV	WSWIGIVYGDGLRSGNVSNELLTNTASSLGFDLKIVNRAPRGSPTVFLSRVYLDPWSSP	233
Redspotted grouper NNV	WSWIGIVYGDGLRSGNVSNELLTNTASSLGFDLKIVNRAPRGSPTVFLSRVYLDPWSSP	233
Japanese flounder NNV	WTWIGIVYGDGLRSGNVSNELLTNTASSLGFDLKIVNRAPRGSPTVFLSRVYLDPWSSP	199
Atlantic halibut NNV	WSWIGIVYGDGLRSGNVSDTLTDTASSLGFDLKIVNRAPRGSPTVFLSRVYLDPWSSP	233
Striped jack NNV	WTWIGIVYGDGLRSGNVSDTLTDTASSLGFDLKIVNRAPRGSPTVFLSRVYLDPWSSP	233
<i>MrNV</i> (West Indies)	FEQTGLSFGDSDLFD-KQYQLRWNYVVEQLGMELKVEFPDPSNG-VTFLARVFPDPYSTN	236
<i>MrNV</i> (China)	FEQTGLSFGDSDLFD-KQYQLRWNYVVEQLGMELKVEFPDPSNG-VTFLARVFPDPYSTN	236
<i>MrNV</i> (India)	FEQTGLSFGDSDLFD-KQYQLRWNYVVEQLGMELKVEFPDPSNG-VTFLARVFPDPYSTN	236
	: * : * * * * . * . :	
<i>Epinephelus</i> NNV	ASVQSPLRLLKLTHTTCDTQSEIDDIGWAKTQAYLVTDCKTPFIGHWCRAQRNCTARVV	293
Sevenband grouper NNV	ASVQSPLRLLKLTHTTCDTQSEIDDIGWAKTQAYLVTDCKTPFIGHWCRAQRNCTARVV	293
Redspotted grouper NNV	ASVQSPLRLLKLTHTTCDTQSEIDDIGWAKTQAYLVTDCKTPFIGHWCRAQRNCTARVV	293
Japanese flounder NNV	-----	293
Atlantic halibut NNV	ASVQSPFRLLKLTHTTCDTQSEIDDIGWAKTQAYLVTDCKTPFIGHWCRAQRNCTARVV	293
Striped jack NNV	ASVQSPLRLLKLTHTTCDTQSEIDDIGWAKTQAYLVTDCKTPFIGHWCRAQRNCTARVV	293
<i>MrNV</i> (West Indies)	TSFQDPLRTRWLNMTSRTCVPEVSAALDRVSGYLVTDKYSPTVSEYCHMIERCYMNAT	296
<i>MrNV</i> (China)	TSFQDPLRTRWLNMTSRTCVPEVSAALDRVSGYLVTDKYSPTVSEYCHMIERCYMNAT	296
<i>MrNV</i> (India)	TSFQDPLRTRWLNMTSRTCVPEVSAALDRVSGYLVTDKYSPTVSEYCHMIERCYMNAT	296

Fig. 3. Multiple sequence alignment of *MrNV* (Indian isolate) with *MrNV* (West Indies and Chinese isolate) and other fishnoda viruses using ClustalW. GenBank accession nos. are *Epinephelus* NNV (AAO73138); Sevenband grouper NNV (AAQ90061); Redspotted grouper NNV(AAQ88211); Japanese flounder NNV (BAB03598); Striped jack NNV (NP_599247); Atlantic halibut NNV(Q9DIC5); *MrNV* (West Indies) (AAO60068); *MrNV* (Chinese) (AAQ54758). NNV represents nervous necrosis virus. The running total number of amino acids is shown on the right. The degree of similarity is illustrated underneath the alignments with a series of consensus symbols. '*' represents residues in that column that are identical in all sequences in the alignment. '.' represents semi conserved substitutions. ':' represents conserved substitutions. Gaps are represented by '-' symbol.

(<http://www.ncbi.nlm.nih.gov/BLAST>) and ClustalW (<http://www.ebi.ac.uk/cgi-bin/clustalW>).

3. Results and discussion

Until recently, disease detection of *MrNV* was based on histological signs and transmission electron microscopy (TEM). A more sensitive detection method by sandwich enzyme linked immunosorbent assay (S-ELISA) was developed by Romestand and Bonami (2003). It was found to have high sensitivity when compared to TEM. However, the availability of antibodies is a major drawback to use this detection method. Much more sensitive and easier genome-based detection methods by dot-blot hybridization, in situ hybridization and RT-PCR for *MrNV* have recently been reported (Widada et al., 2003). Dot-blot and RT-PCR have also been developed for detection of XSV (Widada et al., 2004). In the present study, primers specific to RNA-1 of *MrNV* as reported by Widada et al. (2003) were used (i.e., 1A775 and 1B690) for RT-PCR. This set of primers was designed based on sequence data of the *MrNV* genome (GenBank accession no. AY222839) and was chosen because it had been shown to successfully detect *MrNV* isolates from both the West Indies and China with a sensitivity of 35 fg of RNA template. Using this set of primers reported to amplify a 0.85-kb PCR product, two PCR bands of 850 bp and 750 bp were visualized by agarose gel electrophoresis. The annealing temperature was raised to 58 °C to eliminate the nonspecific 750-bp PCR product, but this modification was not successful. The expected 850-bp PCR product was extracted from the gel and purified. The purified product was then reamplified with the upstream and downstream primers and the resulting PCR product was sequenced (Fig. 1).

The nucleotide sequence obtained from the 850-bp PCR product and its corresponding deduced amino acid sequence are shown in Fig. 2. BLAST analysis of a 738-bp portion of the nucleotide sequence of our *MrNV* Indian isolate showed 98% and 95% identity to nucleotide sequences of *M. rosenbergii* nodavirus, RNA segment-I at GenBank (i.e., accession no. AY222839 and AY231436, respectively). These comparisons confirmed that the PL of Indian *M. rosenbergii* were infected with *MrNV*. The sequence

corresponded to the RNA directed RNA polymerase gene of *MrNV*. Full-length sequencing of Indian *MrNV* is needed for virus strain identification and sequence comparison with other strains isolated from the West Indies and China.

BLAST analysis of 246 deduced amino acid residues of Indian *MrNV* showed very high sequence identity (99%) to those of *M. rosenbergii* nodavirus at GenBank (accession no. AAO60068, AAQ54758). Sequence identity in the range of 33–35% was also observed for the other fish nodaviruses.

ClustalW analysis of 246 deduced amino acid residues of Indian *MrNV* with West Indian and Chinese *MrNV* and other fish nodavirus showed identical, conserved and semiconserved amino acids at various positions (Fig. 3). A total of 60 amino acid residues were found to be identical in all sequences. This sequence conservation is indicative that these viruses are closely related and may have a common ancestor. More detailed genomic analysis and full-length sequencing of Indian *MrNV* are necessary to study the etiology and epidemiology of this disease. Further investigations are also required to find out whether XSV particles are also associated in WTD cases in India.

Acknowledgements

The authors are thankful to Dr. P. Ravichandran, Director, Central Institute of Brackishwater Aquaculture, Chennai, India for the encouragement and facilities provided to carry out this investigation.

References

- Anderson, I.G., Law, A.T., Shariff, M., Nash, G., 1990. A parvo-like virus in the giant freshwater prawn *Macrobrachium rosenbergii*. *J. Invertebr. Pathol.* 55, 447–449.
- Arcier, J.M., Herman, F., Lightner, D.V., Redman, R.M., Mari, J., Bonami, J.R., 1999. A viral disease associated with mortalities in hatchery-reared postlarvae of the giant freshwater prawn *Macrobrachium rosenbergii*. *Dis. Aquat. Org.* 38, 177–181.
- Qian, D., Shi, Z., Zhang, S., Cao, Z., Liu, W., Li, L., Xie, Y., Cambournac, I., Bonami, J.R., 2003. Extra small virus-like particles (XSV) and nodavirus associated with whitish muscle disease in the giant freshwater prawn, *Macrobrachium rosenbergii*. *J. Fish Dis.* 26, 521–527.
- Romestand, B., Bonami, J.R., 2003. A sandwich enzyme linked immunosorbent assay (S-ELISA) for detection of *MrNV* in the

- giant freshwater prawn, *Macrobrachium rosenbergii* (de Man). J. Fish Dis. 26, 71–75.
- Tung, C.W., Wang, C.S., Chen, S.N., 1999. Histological and electron microscopy study *Macrobrachium* muscle virus (MMV) infection in the giant freshwater prawn *Macrobrachium rosenbergii rosenbergii* (de Man), cultured in Taiwan. J. Fish Dis. 22, 1–5.
- Widada, J.S., Durand, S., Cambournac, I., Qian, D., Shi, Z., Dejonghe, E., Richard, V., Bonami, J.R., 2003. Genome based detection methods of *Macrobrachium rosenbergii* nodavirus, a pathogen of the giant freshwater prawn, *Macrobrachium rosenbergii*: dot-blot, in situ hybridization and RT-PCR. J. Fish Dis. 26, 583–590.
- Widada, J.S., Richard, V., Shi, Z., Qian, D., Bonami, J.R., 2004. Dot-blot hybridization and RT-PCR detection of extra small virus (XSV) associated with white tail disease of prawn *Macrobrachium rosenbergii*. Dis. Aquat. Org. 58, 83–87.